

Functional Outcome After Open Reduction Internal Fixation (ORIF) Clavicle Using 1/3 Semitubular Plate

Syed Danish Ali, Khan Shah Azam, Ghazanfar Ali Shah, Syed Muhammad Iqbal, Bassil Azam

ABSTRACT

Objective: To evaluate the functional outcome of ORIF clavicle using DASH score

Methods: This is a retrospective study of prospectively collected data. Patients presented to the Orthopedic Surgery department of TO Clinic from June 2015 till June 2017 with clavicle fracture investigated by clinical and radiological examination who meet the inclusion criteria were treated with open reduction and internal fixation using 1/3 semitubular plate. All patients were operated under general anesthesia. Polysling was applied postoperatively for 4 weeks. Shoulder range of motion was started from 3rd week postoperatively. 6 months after surgery all patients were given a questionnaire to assess their postoperative results using DASH score

Results: 24 patients were enrolled in this paper and more than 90 percent of patients were males. 1 patient didn't complete the study, as he didn't come in the followup. Right clavicle fracture was seen in 20(83.33%) patients while left clavicle fracture was seen in 8(33.33%) patients. The dominating mechanism of injury was fall on shoulder, 22 patients (91.6%). Most of the patients were aged below 40 years, (75%) 18 patients. The average DASH score of patients was 88 (fair to good) .One patients (4.1%) had superficial infection, which resolved with oral antibiotics. Two patients (8.3%) had another fall after surgery, which required redo ORIF. One patient (4.1%) had prominent implant pain, which resolved with pain killers.

Conclusions: We conclude that ORIF Clavicle using 1/3 semitubular plate is a reliable method for clavicle fixation achieving fair to good functional outcome at 6 months after surgery

Keywords: ORIF, 1/3 semitubular plate DASH score

INTRODUCTION

Clavicle is the most frequently injured bone of the body constituting about 2.6% of all fractures [1]. Middle third of the clavicle is mostly vulnerable to fracture reaching upto 70-80% of cases [2]. Management options for clavicle fracture are ORIF, which gives good results especially in displaced fractures [3]. Multiple implants are available for clavicle fixation, plating is considered to be gold standard [4]. The complications of operative intervention include malunion, nonunion, bleeding, infection, and pneumothorax for which

implant removal is being done [5]. Controversy however remains on the best site for plating a clavicle fracture. The two sites commonly preferred for plating are superior and anterior. Both surfaces causes implant prominence and may lead to removal of implant request by the patient [6]. Benefits of anterioinferior plating include usage of long screws [7], avoidance of neurovascular complications and fewer implant related patients complaints [8]. Iatrogenic injury to neurovascular structure can occur regardless of the site of plating [9].

As Pakistan is a third world country where the cost of hospitalization has to be borne by the patient, implant cost is of particular importance. Most of the locking plates available are too costly to be used. The purpose of this study is to evaluate the outcome of using a semitubular plate, which is practically pretty cheap.

METHODS

24 patients from age 18 to 50 yrs. were included in our study with displaced clavicle fracture from June 2015

Department of Orthopaedics

Baqai University Hospital Karachi

Correspondence: Syed Danish Ali

Email: drsyeddanishalia@hotmail.com

Citation: Danish SA, Azam KS, Ghazanfar AS, Iqbal SM, Azam B. Functional Outcome After Open Reduction Internal Fixation (ORIF) Clavicle Using 1/3 Semitubular Plate. J Pak Ortho. 2017 Sep;29(3): page 130-133

till June 2017. All patients had radiologically proven displaced clavicle fracture. Surgery was done under general anaesthesia. A single surgeon having more than 5 years experience did all surgeries. Mostly 6-8 hole 1/3 semitubular plate was applied on the superior surface of the clavicle, wound was closed after securing haemostasis. Poly arm sling was applied postoperatively. Patient was discharged on the 4th postoperative day. Pendulum exercises were started from 2nd week. Stitches were removed after 2 weeks. Physiotherapy was started after 3 weeks. Patients were asked to follow-up at 2nd, 3rd week, 6weeks, 3 months and 6 months postoperatively to assess radiological union. 6 months after surgery all patients were given quick DASH questionnaire [10] to assess surgery results.

Patients with displaced clavicle fracture, Age more than 18 years less than 50yr, Segmental clavicle fracture, Open clavicle fracture and Cosmetic reason were included.

History of previous surgery in the affected side of the shoulder, Ipsilateral shoulder girdle injury, Associated fractures and Diabetes mellitus associated adhesive capsulitis were excluded

Dash Score

Score	Interpretation
98-100	Excellent
93-97	Good to excellent
82-92	Fair to good
66-81	Fair
<= 65	Poor

RESULTS

24 patients were enrolled in this paper and more than 90 percent of patients were males. 1 patient didn't complete the study and was lost in follow up. Right clavicle fracture was seen in 20(83.33%) patients while left clavicle fracture was seen in 8(33.33%) patients.

Table-1: Average age of patient

Age in years	%
> 40	25
< 40	75

Table-3: Mechanism of Injury

Cause of fracture	Number of patients	%
RTA	04	16.66
Fall on shoulder	20	83.33

The dominating mechanism of injury was fall on shoulder, 22 patients (91.6%). Most of the patients were aged below 40 years, (75%) 18 patients. The average DASH score of patients was 88 (fair to good). One patient (4.1%) had superficial infection which resolved with oral antibiotics. Two patients (8.3%) had another fall after surgery, which required redo ORIF.



Figure 1: clavicle fracture



Figure 2: Postoperative x-rays after ORIF clavicle



Figure 3: 3 months postoperative x rays after surgery



Figure 4: Range of motion 3 months after surgery

DISCUSSION

Most of the clavicle fractures unite with conservative methods; commonly employed technique includes figure of eight dressing [11]. None of these conservative methods actually reduce the fracture. Compared to Neers nonunion rate of 1 % recent evidence points to an increased nonunion rate especially in midshaft clavicle fractures when treated conservatively [12]. Nowak et al. showed that in 208 patients 10 years after conservative treatment 96 patients still had symptoms while only 15 patients had nonunion [13]. Anderson in 2003 had showed good functional outcome and low complication rate in 13 patients with lateral clavicle fractures treated with LCP [14]. RCT conducted by the Canadian orthopedic society showed good functional outcome in early plate fixation of displaced midclavicular fractures and decreased nonunion and malunion rates [15]. Another study showed poor functional outcome after conservative treatment of displaced clavicle fractures [16].

There is limited literature available, which shows that 1/3 semitubular plate can be used for clavicle fracture producing good functional outcome with minimum complications. Compared to LCP the cost of 1/3 semitubular plate is minimum so financial restrains can be avoided in our country as most of the important factor in our society for negligence in trauma is financial constrain.

CONCLUSION

In conclusion, we state that ORIF clavicle using 1/3 semitubular plate can be used in selected patients with clavicle fractures producing good results with minimum complication rates.

REFERENCES

1. Neer, C. Fractures of the Clavicle. Fractures in Adults. Rockwood and Green Eds, JB Lippincott, 2nd edition, p 707-713.
2. Brin YS, Palmanovich E, Dolev E, Nyska M, Kish BJ. Displaced mid-shaft clavicular fractures: Is conservative treatment still preferred? *Isr Med Assoc J.* 2014 Dec;16(12):748–52.
3. Rowe CR. An atlas of anatomy and treatment of midclavicular fractures. *Clin Orthop Relat Res.* 1968 May-Jun;58: 29-42.
4. Dhoju D, Shrestha D, Parajuli NP, Shrestha R, Sharma V. Operative Fixation of Displaced Middle Third Clavicl (Edinburg Type 2) Fracture with Superior Reconstruction Plate Osteosynthesis. *Kathmandu Univ Med J (KUMJ).* 2011 Oct-Dec;9(36):286-90.
5. Wenninger JJ, Dannenbaum JH, Branstetter JG, Arrington ED. Comparison of complication rates of intramedullary pin fixation versus plating of midshaft clavicle fractures in an active duty military population. *J Surg Orthop Adv.* 2013;22(1):77–81.
6. Zlowodzki M, Zelle BA, Cole PA, Jeray K, McKee MD. Treatment of acute midshaft clavicle fractures: systematic review of 2144 fractures: on behalf of the Evidence-Based Orthopaedic Trauma Working Group. *J Orthop Trauma* 2005 Aug;19: 504-7.
7. Klein SM, Badman BL, Keating CJ, Devinney DS, Frankle MA, Mighell MA. Results of surgical treatment for unstable distal clavicular fractures. *J Shoulder Elb Surg.* 2010 Oct;19:1049e1055.
8. Böstman O, Manninen M, Pihlajamaki H. Complications of plate fixation in fresh displaced midclavicular fractures. *J Trauma.* 1997 Nov;43:778-783.
9. Collinge C1, Devinney S, Herscovici D, DiPasquale T, Sanders R. Anterior-inferior plate fixation of middle third fractures and nonunions of the clavicle. *J Orthop Trauma.* 2006 Nov-Dec;20:680-686.
10. Jones CB1, Sietsema DL, Ringler JR, Endres TJ, Hoffmann MF. Results of anterior-inferior 2.7mm dynamic compression plate fixation of midshaft clavicular fractures. *J Orthop Trauma.* 2013 Mar;0:1-4.
11. Lo EY, Eastman J, Tseng S, Lee MA, Yoo BJ. Neurovascular risks of anteroinferior clavicular plating. *Orthopedics.* 2010 Jan;33:21.
12. Hudak PL, Amadio PC, Bombardier C. Development of an upper extremity outcome measure: the DASH (disabilities of the arm, shoulder and hand). The Upper Extremity Collaborative Group (UECG) *Am J Ind Med.* 1996 Jun;29(6):602-8.
13. Lester CW. The treatment of fractures of the clavicle. *Ann Surg*1929;89:600-6.
14. Pandya NK, Namdari S, Hosalkar HS. Displaced clavicle fractures in adolescents: Facts,

-
- controversies, and current trends. *J Am Acad Orthop Surg.* 2012 Aug;20(8):498–505.
15. Rehn CH, Kirkegaard M, Viberg B, Larsen MS. Operative versus nonoperative treatment of displaced midshaft clavicle fractures in adults:A systematic review. *Eur J Orthop Surg Traumatol.* 2014 Oct;24(7):1047–53.
 16. Neer CS . Nonunion of the clavicle. *J Am Med Assoc* 1960;172:1006-
 17. McKee MD, Pedesen EM, Jones C, Stephen DJ, Kreder HJ, Schemitsch EH et al. Deficits Following Nonoperative Treatment of Displaced Midshaft Clavicular Fractures. *J Bone Joint Surg (Am)* 2006; 88(1): 35-40.
 18. Anderson K. Evaluation and treatment of distal clavicle fractures. *Clin Sports Med.* 2003;22:319e326.
 19. Nowak J, Holgersson M, Larsson S. Can we predict long-term sequelae after fractures of the clavicle based on initial findings? A prospective study with nine to ten years of follow-up. *J Shoulder Elbow Surg.* 2004 Sep-Oct;13:479-86.
 20. Basamania CJ. Claviculoplasty. *J Shoulder Elbow Surg*, Vol. 8, No. 5, 1999; p 540
 21. Robinson CM, Goudie EB, Murray IR, Jenkins PJ, Ahktar MA, Read EO, et al. Open reduction and plate fixation versus nonoperative treatment for displaced midshaft clavicular fractures: A multicenter, randomized, controlled trial. *J Bone Joint Surg Am.* 2013 Sep;95(17):1576–84
 22. McKee RC, Whelan DB, Schemitsch EH, McKee MD. Operative versus nonoperative care of displaced midshaft clavicular fractures: A meta-analysis of randomized clinical trials. *J Bone Joint Surg Am.* 2012 Apr;94(8):675–84.